

Charge Symmetry Violation in $pn \rightarrow d\pi^0$ as a Test of Chiral Effective Field Theory

U. van Kolck ^{a,b,1}, J.A. Niskanen ^{c,2}, and G.A. Miller ^{b,3}

^a *Kellogg Radiation Laboratory, 106-38, California Institute of Technology,
Pasadena, CA 91125, USA*

^b *Department of Physics, University of Washington,
Seattle, WA 98195, USA*

^c *Department of Physics, University of Helsinki,
FIN-00014 Helsinki, Finland*

Abstract

Chiral effective field theory predicts a specific charge symmetry violating amplitude for pion production. This term is shown to provide the dominant contribution to the forward-backward asymmetry in the angular distribution for the reaction $pn \rightarrow d\pi^0$, for reasonable values of the mass difference between down and up quarks, δm_N . Using a value $\delta m_N \approx 3$ MeV leads to a prediction of a 10 standard deviation effect for a TRIUMF experiment.

¹vankolck@krl.caltech.edu

²jouni.niskanen@helsinki.fi

³miller@phys.washington.edu

Charge symmetry is an approximate symmetry of QCD. If one ignores electromagnetic corrections and the mass difference between u and d quarks, then the Lagrangian is invariant under the interchange of $u \leftrightarrow d$. Isospin invariance implies charge symmetry, but the converse is not true.

Much has been learned about charge symmetry breaking (CSB); see the reviews [1, 2]. The general importance of the quark mass difference effect was stressed in Ref. [2]. Our purpose here is to use chiral effective field theory to make predictions, based on QCD, for the reaction $np \rightarrow d\pi^0$.

The use of chiral Lagrangians, which employ hadronic degrees of freedom, represents a serious effort to provide a rigorous and model-independent methodology to use QCD to make predictions at relatively low energies [3, 4, 5]. This is because the most general Lagrangian which respects unitarity, has correct properties under cluster decomposition and the same symmetries as QCD, should give results that encompass the results from QCD proper [6]. Predictive power is retained because at low energies an expansion in momentum can be formulated using power counting arguments.

We wish to exploit the feature that chiral symmetry makes predictions for effects that stem from the small but *explicit* breaking of chiral symmetry generated by the quark masses. Consider the proton-neutron mass difference $m_n - m_p$, which includes a contribution δm_N from the quark mass difference $m_d - m_u \equiv \varepsilon(m_d + m_u)$ ($\varepsilon \sim 1/3$), and another contribution $\bar{\delta} m_N$ from electromagnetic effects. Thus $m_n - m_p = \delta m_N + \bar{\delta} m_N$. Neglecting strangeness, chiral symmetry is essentially an $SO(4)$ internal symmetry. One can show that the quark-mass-difference term in the QCD Lagrangian behaves under $SO(4)$ as the third component of an $SO(4)$ vector [7, 8, 9]. Therefore, in the effective hadronic theory, all isospin-violating interactions generated by the quark mass difference break $SO(4)$ as third components of (tensor products of) $SO(4)$ vectors. The operators of interest here involve the nucleon N and pion π fields. The leading isospin-violating term at low energies coming from the quark mass difference is the third component of an $SO(4)$ vector [8, 9]:

$$\mathcal{L}_{\text{qm}}^{(1)} = \frac{\delta m_N}{2} \left(N^\dagger \tau_3 N - \frac{2}{DF_\pi^2} N^\dagger \pi_3 \boldsymbol{\pi} \cdot \boldsymbol{\tau} N \right), \quad (1)$$

where $\boldsymbol{\tau}$ represents the Pauli matrices in isospin space, $F_\pi = 186$ MeV is the pion decay constant, and $D = 1 + \boldsymbol{\pi}^2/F_\pi^2$. We use $D = 1$ here. Eq. (1) represents the isospin-violating part of the nucleon sigma term.

Furthermore, one can show that quark interactions generated by (“hard”) photon exchange break $SO(4)$ as the 34 component of an $SO(4)$ antisymmetric rank-2 tensor [8]. Thus the low-energy effective theory must include isospin-violating interactions that break $SO(4)$ as 34 components of (tensor products of) $SO(4)$ antisymmetric rank-2 tensors. The leading isospin-violating term of this kind at low energies was shown to be [8]:

$$\mathcal{L}_{\text{hp}}^{(-1)} = \frac{\bar{\delta} m_N}{2} \left(N^\dagger \tau_3 N + \frac{2}{DF_\pi^2} N^\dagger (\pi_3 \boldsymbol{\pi} \cdot \boldsymbol{\tau} - \boldsymbol{\pi}^2 \tau_3) N \right). \quad (2)$$

Observe that chiral symmetry links the first terms in Eqs. (1,2), which are contributions to the nucleon mass difference, to the second terms, which are isospin-violating

pion-nucleon interactions. Since the parameter F_π is determined from another process (pion decay), these pion-nucleon interactions *must exist* with the given strength if our understanding of QCD is not totally flawed.

Let us discuss the values of the parameters, δm_N and $\bar{\delta} m_N$, which are not determined by chiral symmetry. The sum of these terms is fixed by the nucleon mass difference $m_n - m_p = 1.3$ MeV. Dimensional analysis suggests their separate order of magnitude in terms of the natural QCD mass scale, $M_{QCD} \sim 1$ GeV. We know $\delta m_N \propto m_d - m_u$ and $m_\pi^2 \propto m_d + m_u$, so $\delta m_N = O(\varepsilon m_\pi^2 / M_{QCD}) \sim 7$ MeV. The electromagnetic term $\bar{\delta} m_N$, long known to be negative [10], should be proportional to the fine structure constant α , so that $\bar{\delta} m_N = -O(\alpha M_{QCD} / \pi) \sim -2$ MeV. Estimates such as these cannot be trusted to better than a factor of a few. More precise estimates can only be made at present with model-dependent assumptions. Most models estimate that $\bar{\delta} m_N \approx -\alpha / R_N \simeq -1.5$ MeV [2], in which case $\delta m_N \simeq 3$ MeV. We will refer to these values, which are also consistent with the results of Gasser and Leutwyler [11], as “quark model estimates”.

Verifying the existence of the terms of Eqs. (1,2) is of high interest, as is determining the parameters. The simplest procedure would be to study pion-nucleon scattering close to threshold [7, 8, 9, 12]. The lack of π^0 beams requires the comparison of charge exchange and elastic scattering processes. If one neglects electromagnetic effects and the kinematic dependence on the threshold values of the energy on the masses, the leading-order chiral Lagrangian gives for the triangle discrepancy among the pion-nucleon amplitudes T :

$$D = -2 \frac{T(\pi^+ p \rightarrow \pi^+ p) - T(\pi^- p \rightarrow \pi^- p) - \sqrt{2} T(\pi^- p \rightarrow \pi^0 n)}{T(\pi^+ p \rightarrow \pi^+ p) - T(\pi^- p \rightarrow \pi^- p) + \sqrt{2} T(\pi^- p \rightarrow \pi^0 n)} \approx - \frac{\delta m_N - \bar{\delta} m_N}{2m_\pi}. \quad (3)$$

The quantity D is found [13] to be about 6%, leading to a very large **negative** value of $\delta m_N - \bar{\delta} m_N \approx -17$ MeV. If the 6% result is correct, there seems to be a serious problem with our current understanding of low-energy QCD. However, electromagnetic effects are very important at low energies as are the kinematic difference in threshold energies. Kinematic effects and sub-leading strong interactions have been studied in Ref. [12], but a precise accounting for the electromagnetic interactions presents a very serious challenge to theorists. A small shift in the normalization could be reflected in a huge change in D . It seems that a successful determination of δm_N and $\bar{\delta} m_N$ from πN scattering will require a strong theoretical and experimental effort.

The biggest problem in all of this is the lack of a pion beam. The reaction $np \rightarrow d\pi^0$ is relevant because it involves the production of a virtual pion from one nucleon (“pion bremsstrahlung”), followed by rescattering on the second. Charge symmetry predicts that the angular distribution is symmetric about 90° in the center-of-mass. Any asymmetry must be caused by charge-symmetry-violating effects. We here examine the integrated forward-backward asymmetry in the center of mass of the reaction $np \rightarrow d\pi^0$,

$$A_{fb} = \frac{\int_0^{\frac{\pi}{2}} d\Omega [\sigma(\theta) - \sigma(\pi - \theta)]}{\int_0^\pi d\Omega \sigma(\theta)}. \quad (4)$$

Our interest in this particular observable is motivated by an experiment in progress at

TRIUMF [14] which aims to measure this asymmetry close to threshold, $E_{lab} = 279.5$ MeV or $\eta = q_\pi/m_\pi = 0.17$, with a precision of $\pm 0.06\%$ ⁴.

A_{fb} arises from an interference between s - and p -wave pions and thus decreases as one approaches threshold energies. Nevertheless, there are advantages to considering this pion production reaction at threshold kinematics. Near threshold one minimizes the four-momentum transferred k , enhancing isospin violation relative to chiral invariance, because chiral (thus isospin) conserving contributions to πN rescattering vanish with the pions' four-momenta. Right at threshold the leading πN rescattering mechanism (Weinberg-Tomozawa term [15]) goes as the sum of the pions' energies, $\omega_q + \omega_k$; the real pion has as small energy as possible, $\omega_q \simeq m_\pi$, while the typical energy of the virtual pion is just $\omega_k \sim m_\pi/2$. In pion production the three-momentum $p \approx \sqrt{m_N m_\pi}$ of the virtual pion is relatively large, which slows the convergence of a momentum expansion. However, there is a trade-off here, since the large momentum reduces the importance of electromagnetic effects. Note, also, that at any given order in a momentum expansion in p/M_{QCD} , the number of relevant isospin-breaking contributions is more limited than isospin-conserving ones. This leads to an isolation of δm_N and $\bar{\delta} m_N$ effects. Even if other mechanisms (*e.g.* π - η mixing) are also relevant, they are not expected to exactly cancel among themselves. Thus, a model adjusted to correctly describe the total cross-section might provide a reliable estimate of the overall size of the δm_N and $\bar{\delta} m_N$ contributions to the asymmetry; and these estimates would be confronted with data.

The most recent calculation of the near-threshold asymmetry [16] includes effects of the nucleon-mass splitting on the πN vertex function, which influences both the pion production amplitude and the nucleon-nucleon potential, and the far larger influence of π - η - η' mixing, both in the pion-production kernel and in initial and final interactions between nucleons. This calculation does not include the seagull interactions (1,2) proportional to δm_N and $\bar{\delta} m_N$. The isospin-conserving interactions are included in a nucleon-delta coupled-channel model that describes both pion production and NN phase shifts to an accuracy of a few degrees from threshold over the delta region [17]. Close to threshold π - η mixing is the dominant effect, taking place both in pion s and p waves.

Here we estimate, using both a power counting argument and a numerical evaluation, the effect of δm_N and $\bar{\delta} m_N$ in the asymmetry (4) close to threshold. These mechanisms appear in the s wave, and can be relatively large because the isospin-conserving p wave amplitude is not vanishingly small when compared to the isospin-conserving s wave, even below 300 MeV. A power counting argument suggests immediately that the contributions from Eqs. (1,2) should be dominant. Let us take the contribution from the δm_N seagull, Fig. 1a, as an example. The isospin-conserving p wave emission is proportional to q/F_π , while the isospin-violating s wave gets a p/F_π from the virtual pion emission, $1/p^2$ from the pion propagator, and $\delta m_N/F_\pi^2$ from the seagull vertex. The product of the two amplitudes is then expected to be

$$\mathcal{M}_{qm} \sim \varepsilon \left(\frac{m_\pi}{m_N} \right)^{3/2} \frac{q}{F_\pi^4}, \quad (5)$$

⁴The definition of the asymmetry A_{fb} used here is 1/2 of the definition used in Ref. [14].




Figure 1: Contributions to A_{fb} : (a) $\mathcal{M}_{qm}$; (b) $\mathcal{M}_{\pi\eta}$. Here nucleon (pion) propagation is represented by a solid (dashed) line; and pion emission/absorption is denoted by a (circled) solid dot in case of an isospin-conserving p (s) wave, and a (circled) cross in case of an isospin-violating s (p) wave.

in which the notation qm is used to denote the effects of Eq. (1). On the other hand, the contribution from π - η mixing is smaller by a factor of m_π/m_N near threshold. This can be seen by integrating out the η . This produces an isospin-violating pion-nucleon coupling of size $\beta q/F_\pi$ in the p wave and $\beta\omega_q p/F_\pi m_N$ in the s wave, where $\beta \sim \varepsilon m_\pi^2/m_N^2$ [19]. Let us consider now isospin violation in the p -wave pion emission, Fig. 1b. The isospin-conserving s -wave emission is proportional to $\omega_q p/F_\pi m_N$, while the isospin-violating p wave gets a p/F_π from the virtual pion emission, $1/p^2$ from the pion propagator, another p/F_π from the virtual pion absorption, $1/\omega_q$ from the nucleon propagator, and $\beta q/F_\pi$ from the real pion emission. The product of the two amplitudes is then expected to be

$$\mathcal{M}_{\pi\eta} \sim \varepsilon \left(\frac{m_\pi}{m_N} \right)^{5/2} \frac{q}{F_\pi^4}. \quad (6)$$

The same estimate holds for the interference between isospin-conserving p wave and isospin-violating s wave. A comparison of Eqs. (5) and (6) reveals the ratio m_π/m_N between the two effects. Note that the counting rules used here are consistent with those of previous works in pion production[18].

We calculate A_{fb} following the procedure described in Ref. [16] and earlier references. Before presenting the results, we display aspects of the relevant operators. Denoting the two nucleons with superscripts (1) and (2) , the spin-isospin structure of the isospin-violating rescattering is (at threshold, and neglecting terms of order m_π/m_N)

$$\begin{aligned} \mathcal{O}_{qm} = & -\frac{f}{4\pi\mu F_\pi^2 \sqrt{2}\mu} [\delta m_N (\boldsymbol{\tau}^{(1)} \cdot \boldsymbol{\tau}^{(2)} + \tau_3^{(1)} \tau_3^{(2)}) - \bar{\delta} m_N (\boldsymbol{\tau}^{(1)} \cdot \boldsymbol{\tau}^{(2)} - \tau_3^{(1)} \tau_3^{(2)})] \\ & (\boldsymbol{\sigma}^{(1)} - \boldsymbol{\sigma}^{(2)}) \cdot (-i\hat{\mathbf{r}}) \frac{d}{dr} \left(\frac{e^{-\mu' r}}{r} \right), \end{aligned} \quad (7)$$

where μ is the charged pion mass, $\mu' = \sqrt{3}\mu/2$, and f is the pion-nucleon coupling constant ($f = g_A \mu/F_\pi$ in leading order; $f^2/4\pi = 0.075$ is used here). The isospin operator allows transitions from states with isospin $I = 0$ to the deuteron, also $I = 0$:

$$\langle I = 0 | \boldsymbol{\tau}^{(1)} \cdot \boldsymbol{\tau}^{(2)} \pm \tau_3^{(1)} \tau_3^{(2)} | I = 0 \rangle = -3 \mp 1, \quad (8)$$

where only the isospin is indicated. Thus there is a large matrix element. If we do the spin-angle evaluation of the matrix element of $\mathcal{O}_{\text{qm}}$ between 1P_1 and 3S_1 (for the deuteron) states we find

$$\langle ^3S_1 | \mathcal{O}_{\text{qm}}(r) | ^1P_1 \rangle = \frac{2f}{\pi\mu F_\pi^2 \sqrt{3}\sqrt{2\mu}} (\delta m_N - \frac{\bar{\delta}m_N}{2}) (-i) \frac{d}{dr} \left(\frac{e^{-\mu' r}}{r} \right). \quad (9)$$

We emphasize that the asymmetry in pion production thus depends on a different combination of δm_N and $\bar{\delta}m_N$ than either the total nucleon mass splitting or the isospin violation related to charge-exchange pion-nucleon scattering. Under the assumption that we have the isospin-conserving mechanisms of pion production under control, A_{fb} can be used in conjunction with $m_n - m_p$ to extract δm_N and $\bar{\delta}m_N$.

The above operator may be compared with the operator involving phenomenological isospin-conserving, on-shell pion rescattering, which in leading order is given by the Weinberg-Tomozawa term [15], sometimes written in terms of $\lambda_2 = \mu^2/4\pi F_\pi^2 \approx 0.045$. If we consider threshold kinematics, and only the matrix element between 3P_1 and 3S_1 states, we obtain [20]

$$\mathcal{O}_{\text{ic}} = -\frac{3f}{2\mu^2\sqrt{2\mu}} \lambda_2 i(\boldsymbol{\tau}^{(1)} \times \boldsymbol{\tau}^{(2)})_3 (\boldsymbol{\sigma}^{(1)} + \boldsymbol{\sigma}^{(2)}) \cdot (-i\hat{\mathbf{r}}) \frac{d}{dr} \left(\frac{e^{-\mu' r}}{r} \right), \quad (10)$$

so that

$$\langle ^3S_1 | \mathcal{O}_{\text{ic}}(r) | ^3P_1 \rangle = 6\sqrt{\frac{2}{3}} \frac{f}{\mu^2\sqrt{2\mu}} \lambda_2 (-i) \frac{d}{dr} \left(\frac{e^{-\mu' r}}{r} \right). \quad (11)$$

Assuming that the initial np 3P_1 and 1P_1 wave functions are the same (the phase shifts differ by only $\sim 10\%$ at our energies) so that the radial matrix elements of the same, we obtain a rough estimate for the ratio of matrix elements:

$$\frac{\langle ^3S_1 | \mathcal{O}_{\text{qm}} | ^1P_1 \rangle}{\langle ^3S_1 | \mathcal{O}_{\text{ic}} | ^3P_1 \rangle} \approx \frac{2\sqrt{2}}{3} \frac{\delta m_N - \frac{\bar{\delta}m_N}{2}}{m_\pi} \approx 0.03, \quad (12)$$

using the quark model estimate $\delta m_N - \bar{\delta}m_N/2 \simeq 4$ MeV. This comes from taking $\delta m_N = 3.1$ MeV[2] so that $\bar{\delta}m_N = -1.8$ MeV. This provides a first estimate of A_{fb} in the percent range. (Cf. Eq. (3) for the triangle discrepancy.)

We turn now to the results of our detailed numerical evaluations. For illustration here we again use the quark model estimate $\delta m_N - \bar{\delta}m_N/2 \simeq 4$ MeV. The amplitudes arising from using this value in Eq. (7) (denoted as loosely as “qm”), as well as those obtained from the sum of the previously computed amplitudes of Ref.[16] (denoted loosely as $\pi\eta$), are listed in Table 1. These are reduced matrix elements in the definition used *e.g.* in Ref. [21]. Some overall kinematic factors are ignored.

At 279.5 MeV, the energy of the TRIUMF experiment, the computed contribution from δm_N and $\bar{\delta}m_N$ to the asymmetry of Eq. (4) is found to be 0.97%, or about -3.5 times the strength of (and opposite in sign to) the sum of the contributions calculated in Ref. [16]. This earlier calculation gave $A_{fb} = -0.28\%$. As a consequence, the *total* asymmetry is

Table 1: Real and imaginary parts of amplitudes at $\eta = 0.17$, $E = 279.5$ MeV. The CSB amplitudes have been multiplied by a factor of 10^3 . See the text for the definition of qm and $\eta\pi$.

transition	qm amplitude		$\pi\eta$ amplitude	
	real part	imag part	real part	imag part
$^1P_1 \rightarrow s$	25.06	-13.25	-10.38	6.34
$^3S_1 \rightarrow p$	3.01	0.68	-3.86	-1.36
$^3D_1 \rightarrow p$	0.82	-0.10	-7.26	2.65
$^3D_2 \rightarrow p$	6.47	2.76	2.89	-0.3
$^3P_1 \rightarrow s$	0.901	-0.487	0.901	-0.487
$^1D_2 \rightarrow p$	0.535	0.086	0.535	0.086

$A_{fb} = 0.69\%$, three times the asymmetry of Ref. [16], with opposite sign. This prediction is equivalent to a 10 standard deviation effect for the experiment of Ref. [14].

While such a large result is anticipated from the estimates given above, it is worthwhile to provide an analysis of the amplitudes. These results can be qualitatively understood by looking at the interference of the most important low partial wave amplitudes (denoted in the spectroscopic notation by $l_{\pi J}$ or $l_{\pi LJ}$; the hat indicates CSB amplitudes)

$$\sigma(\theta) - \sigma(\pi - \theta) = 4\sqrt{\frac{2}{3}} \text{Re } p_2^* \hat{s}_1 \cos \theta + 2\sqrt{2} \text{Re } s_{11}^* (\hat{p}_{22} - \sqrt{\frac{2}{3}} \hat{p}_{01} + \sqrt{\frac{1}{3}} \hat{p}_{21}) \cos \theta. \quad (13)$$

This can be compared with the charge-symmetric cross section

$$\sigma(\theta) + \sigma(\pi - \theta) = 2|s_{11}|^2 + \frac{1}{3}|p_2|^2 + |p_2|^2 \cos^2 \theta. \quad (14)$$

(The small amplitude $^1S_0 \rightarrow p_0$ can be neglected in this qualitative comparison.) For the case of qm, the largest two CSB amplitudes dominate so that one can clearly see that the two interfering terms above add constructively. The other CSB terms, computed earlier, have different phases and are therefore subject to destructive interference.

Note that the total cross section computed from the present model at threshold yields an overestimate of the data by about 50% [23]. We discuss how this could influence our present result. The amplitude s_{11} is dominant, so one could arbitrarily reproduce the data for the total cross section by dividing the amplitude by $\sqrt{1.5}$. This amplitude enters in the second (smaller) term of Eq. (13) and in the denominator, so that the resulting value of A_{fb} (for all contributions) would increase by a factor of about 1.4. However, the same pion rescattering mechanism appears in the both charge symmetric and CSB amplitudes. In that case, the error could cancel out in the ratio A_{fb} . We take an error estimate from the geometric mean of unity and 1.4, which gives a theoretical uncertainty of about 20%

As the isospin-violating amplitudes are small, they are approximately linear in $\delta m_N - \bar{\delta} m_N/2$. Thus we are able to parametrize A_{fb} as a linear function of $\delta m_N - \bar{\delta} m_N/2$, at

any energy. At $E = 279.5$ MeV, using the previous results we write

$$A_{fb} \simeq -.28\% \times \left(1 - \frac{0.87}{\text{MeV}} (\delta m_N - \frac{\bar{\delta} m_N}{2}) \right). \quad (15)$$

The predicted value of A_{fb} depends strongly on the value of quantity $\delta m_N - \bar{\delta} m_N/2$. Indeed, A_{fb} nearly vanishes if this is 1 MeV instead of 4 MeV. However, there are estimates cited in the reviews [1, 2, 11] which would obtain $\delta m_N - \bar{\delta} m_N/2$ significantly greater than 4 MeV.

The energy dependence of the amplitudes around this energy is as expected (s wave independent of momentum, p wave proportional to q). The asymmetry is not quite as good in its energy dependence as the elementary amplitudes: below 300 MeV it is reasonably proportional to q ; however, above this energy it begins to curve down. With the quark model estimate, A_{fb} increases to 1.76% at 300 MeV and 1.87% at 320 MeV.

The principal conclusion of the present paper, based on Eq. (15) is that the successful measurement of the value of the forward-backward asymmetry of Eq. (4) for the $np \rightarrow d\pi^0$ reaction will provide a serious test of our understanding of QCD in terms of chiral effective field theory. We look forward to the results of Ref. [14] with great anticipation.

We close the paper by suggesting that other experiments probing isospin violation could supply additional information, in the whole providing an even more stringent test. We have already mentioned the need for more πN data. Another experiment [22], planned at IUCF, would measure the cross section for $dd \rightarrow \alpha\pi^0$. Charge symmetry prevents the reaction $dd \rightarrow \alpha\pi^0$ from occurring, so that any non-zero cross section must be due to the breaking of the symmetry. The cross section for $dd \rightarrow \alpha\pi^0$ is proportional to the square of the matrix element of the CSB pion-production operator. Thus the cross section is very small and hard to measure. However, there is no interference with amplitudes that respect charge symmetry, which would simplify the interpretation of any measured cross section. Furthermore, purely electromagnetic effects are very small, and symmetries—parity, angular momentum conservation and the identical nature of the two initial deuterons— forbid the production of a pion in a p wave. These features simplify the analysis of the reaction $dd \rightarrow \alpha\pi^0$.

Acknowledgments

This research was supported in part by the Academy of Finland and the Finnish Academy of Science and Letters through the Vilho, Yrjö and Kalle Väisälä Foundation, the U.S. Department of Energy, and the U.S. National Science Foundation. We thank C. Hanhart for useful comments. Two of us (G.A.M and J.A.N.) thank ECT* where this work was completed.

References

- [1] E.M. Henley, in “Isospin in Nuclear Physics”, ed. D.H. Wilkinson, North-Holland, Amsterdam, 1969; E.M. Henley and G.A. Miller, in “Mesons in Nuclei, Vol. I”, ed. M. Rho and D.H. Wilkinson, North-Holland, Amsterdam, 1979; G.A. Miller and W.T. van Oers, in “Symmetries and Fundamental Interactions in Nuclei”, ed. W.C. Haxton and E.M. Henley, World Scientific, Singapore, 1995.
- [2] G.A. Miller, B.M.K. Nefkens, and I. Šlaus, Phys. Rep. **194** (1990) 1.
- [3] “Chiral Dynamics: Theory and Experiment”, ed. A.M. Bernstein and B.R. Holstein, Springer, Berlin, 1995.
- [4] “Nuclear Physics with Effective Field Theory II”, ed. P. Bedaque, M.J. Savage, R. Seki, and U. van Kolck, World Scientific, Singapore, 1998.
- [5] U. van Kolck, Prog. Part. Nucl. Phys. **43** (1999) 337.
- [6] S. Weinberg, Physica **96A** (1979) 327.
- [7] S. Weinberg, Trans. N.Y. Acad. Sci. **38** (1977) 185.
- [8] U. van Kolck, Ph.D. dissertation, U. of Texas (1993); Few-Body Syst. Suppl. **9** (1995) 444.
- [9] S. Weinberg, in Ref. [3], p. 3.
- [10] A. Zee, Phys. Rep. **3** (1972) 127.
- [11] J. Gasser and H. Leutwyler, Phys. Rep. **87** (1982) 77.
- [12] N. Fettes, U.-G. Meißner, and S. Steininger, Phys. Lett. **B451** (1999) 233.
- [13] W.R. Gibbs, L. Ai, and W.B. Kaufmann, Phys. Rev. Lett. **74** (1995) 3740; E. Matsinos, Phys. Rev. C **56** (1997) 3014.
- [14] A.K. Opper and E. Korkmaz (spokespersons), TRIUMF E-704 Proposal.
- [15] S. Weinberg, Phys. Rev. Lett. **17** (1966) 616; Y. Tomozawa, Nuovo Cim. **46A** (1966) 707.
- [16] J.A. Niskanen, Few-Body Syst. **26** (1999) 241.
- [17] J.A. Niskanen, Nucl. Phys. **A298** (1978) 417; Phys. Rev. C **45** (1992) 2648.
- [18] T. D. Cohen, J.L. Friar, G. A. Miller, and U. van Kolck, Phys.Rev. C **C53**, 2661 (1996); C. Hanhart, U. van Kolck, and G.A. Miller, nucl-th/0004033
- [19] U. van Kolck, J.L. Friar, and T. Goldman, Phys. Lett. **B371** (1996) 169.
- [20] D.S. Koltun and A. Reitan, Phys. Rev. **141** (1966) 1413.

- [21] A. deShalit and I. Talmi, “Nuclear Shell Theory”, Academic Press, New York, 1969.
- [22] A.D. Bacher and E. Stephenson (spokespersons), IUCF CE-78 Proposal.
- [23] J.A. Niskanen, Phys. Rev. C **53** (1996) 526.